

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**INSTRUCTIONS FOR USE****PPP: Ceta SG (PCS No. 06252)**

CROPS: *Forestry and Forest Nursery*

For this EAMU the following crop definitions apply:

*Forest nursery is considered as outside cultivation only.

This approval provides for the use of the product **Ceta SG (PCS No. 06252)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 29.01.19. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Ceta SG
<i>Active Substance:</i>	Acetamiprid
<i>Formulation:</i>	20% w/w (Water Soluble Granules)
<i>Authorisation Holder:</i>	Unichem Ltd
<i>Marketing Company:</i>	Unichem Ltd
<i>PCS No:</i>	06252
<i>User type/Function:</i>	Professional Use Only / Insecticide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

<i>Crops:</i>	<i>Maximum Individual Dose:</i>	<i>Max No of applications:</i>	<i>Maximum total dose:</i>	<i>Harvest Interval :</i>	<i>Target:</i>
Forest nursery	250 g/ha	1 per year	250 g/ha	N/A	Large Pine weevil (<i>Hylobius spp.</i>)
Forestry	*Pre planting	1	0.185g/plant		Large Pine weevil (<i>Hylobius spp.</i>)
	**Post planting	1	0.185g/plant		Large Pine weevil (<i>Hylobius spp.</i>)

Restrictions:

**Apply at a rate of 0.185 g/plant to be applied in a minimum water volume of 1.5 mL/plant through hydraulic nozzles, spinning disc, electrodyne electrostatic applicator in an enclosed cabinet or dipping roots and shoulder in a dipping module at the rate indicated above.*

***Plants treated prior to planting may have a follow up treatment applied between March and October in the year of planting. Apply at a rate of 0.185 g/plant in a minimum water volume of 20mL/plant.*

*Maximum amount applied shall not exceed **832.5 g/ha**.*

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient(s).
- (3) Forestry (Pre Planting): Enclosed cabinet via hydraulic nozzles, spinning disc or electrodyne electrostatic application or via automated dipping module.
- (4) Forestry (Post Planting): Knapsack/hand held sprayers only.
- (5) Do not apply more than 0.8325 kg/ha of Ceta SG to the same area of forestry land per annum.
- (6) After Pre Planting application, do not handle until transplants are dry.
- (7) After application do not enter treated areas until the treated area is completely dry.
- (8) This approval extends to professional use only.
- (9) For operator protection refer to the product label in conjunction with these instructions.
- (10) For environmental information refer to the product label in conjunction with these restrictions but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies from knapsack sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (11) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

This approval relates to use of “Ceta SG” as a horticultural insecticide for use on the named crops and situations listed above.

Ceta SG is an insecticide with systemic activity.

Resistance Management:

Apply Ceta SG as a protectant spray or at the first sign of pest attack.

Use “Ceta SG” as part of an Integrated Crop Management (ICM) strategy.

Ceta SG contains one active ingredient classed within the group: Neonicotinoid (IRAC 4A). Use no more than two applications of any neonicotinoid insecticide on any crop. Where a neonicotinoid insecticide has previously been used on the crop (e.g., prior to planting) use no more than a single foliar application of a neonicotinoid.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical types or alternative control measures should be included in the planned programme. Alternating insecticides with different modes of action is a recognised anti-resistance strategy and Ceta should always be used in conjunction with other insecticides with a different mode of action. This includes consideration of sequences involving seed treatments followed by foliar sprays. Where a neonicotinoid seed has been used, the first subsequent foliar spray should be a non-neonicotinoid product with a different mode of action / different cross resistance group with satisfactory efficacy against the targeted pest.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.